

SPINNING PARTICLES IN 2+1 DIMENSIONS

Subir Ghosh

Department of Physics

Gobardanga Hindu College, 24 Pgs. (North)

West Bengal, India

Abstract: Lagrangian and Hamiltonian formulations of a free spinning particle in 2+1-dimensions or *anyon* are established, following closely the analysis of Hanson and Regge. Two viable (and inequivalent) Lagrangians are derived. It is also argued that one of them is more favourable. In the Hamiltonian analysis non-trivial Dirac Brackets of the fundamental variables are computed for both the models. Important qualitative differences with a recently proposed model for anyons are pointed out.

Particles in 2+1-dimensions are allowed to possess arbitrary spin and statistics. These so called anyons have been attracting a great deal of attention [1] due to their varied applications in planar physics, eg. fractional quantum Hall effect, high T_c superconductivity and in processes in the presence of cosmic strings. In our classical analysis as well, we will refer to the spinning particles as anyons.

In order to proceed to the field theory of anyons, it becomes imperative to study the free anyon. An important step in this direction is the work by Jackiw and Nair [2], who have derived an equation for the anyon, (in analogy to the Dirac equation for spin $\frac{1}{2}$ particle). Starting from a Poincare group representation point of view, they have taken the four-momentum square and the Pauli -Lubanski scalar (since the space time is 2+1-dimensional) as the *two natural independent* choices of invariants. This is important for our formulation as well since we will consider the model which has the above characteristics, as the more favourable one. On the other hand, a covariant Lagrangian description of anyon has been attempted by Chaichian, Gonzalez Felipe and Martinez [3]. Also in [3], a Hamiltonian constraint analysis [4] has been carried through. We have expressed our reservations regarding this model at the end of the Letter.

In the present Letter, a systematic way of deriving the anyon Lagrangian has been presented. This was formulated by Hanson and Regge [5] much earlier, for the case of spinning particles in 3+1-dimensions. However, a detailed analysis reveals that there are interesting differences between 3+1 and 2+1-dimensional models, even at the classical level. As we will elaborate, the formulation is based on deriving a consistency condition, involving the Lagrangian L , the squared linear and angular momentum and the basic degrees of freedom of the anyon. Allowed expressions for L ought to come out as solutions of the consistency equation. Unfortunately this equation is too complicated to solve and an eiconal procedure is adopted instead [5] in deriving one particular explicit expression for an L . This model obeys the famous Regge Trajectory (RT), that connects its mass and spin value. On the other hand, consistent with this scheme, another type of L is also

possible which *does not obey the RT*. Indeed, this was already noted in [5] under some special circumstances. We argue that the latter model is more preferable, since it also agrees with the basic assumptions of the Jackiw-Nair model [2]. Let us now move on to the actual analysis.

Following Hanson and Regge [5], the anyon is depicted as a point on the world line, to which a rotating frame is attached. The (tentative) Lagrangian coordinates are taken as the position $x^\mu(\tau)$, and angle variables $\Lambda_\nu^\mu(\tau)$, with the restriction

$$\Lambda^\mu{}_\lambda \Lambda^{\nu\lambda} = \Lambda_\lambda{}^\mu \Lambda^{\lambda\nu} = g^{\mu\nu}; g^{00} = -g^{11} = -g^{22} = 1 \quad (1)$$

where τ fixes the position on the world line. The velocities are

$$\dot{x}^\mu(\tau) = \frac{dx^\mu(\tau)}{d\tau} \equiv u^\mu(\tau); \dot{\Lambda}^\mu{}_\nu(\tau) = \frac{d\Lambda^\mu{}_\nu(\tau)}{d\tau}. \quad (2)$$

From (1) we find

$$\Lambda_\lambda{}^\mu \dot{\Lambda}^{\lambda\nu} = \sigma^{\mu\nu} = -\sigma^{\nu\mu}. \quad (3)$$

Thus $\sigma^{\mu\nu}$ happens to be a better analogue of the usual angular velocity (instead of $\dot{\Lambda}_\nu{}^\mu$). Note the $\sigma^{\mu\nu}$ contains more than the requisite number of components since in two spatial dimensions, spin has a single component. Eventually we will remove the extra degrees of freedom by imposing constraints. We can still restrict the various combinations of u^μ and $\sigma^{\mu\nu}$ such that a Poincare invariant Lagrangian is obtained. This calls for a subtle analysis [5] since *two* reference frames are involved here, *i.e.* the (laboratory) fixed frame and the one fixed on the anyon. In the anyon frame L will be a Lorentz scalar. However, Poincare invariance demands that L should not be affected by translations and rotations of the fixed frame. Thus the variables comprising L should be invariant (even though they may not be Lorentz scalars), with respect to transformations of the fixed frame. The above analysis restricts the Lagrangian variables to only three scalar combinations

$$a_1 = u^\mu u_\mu, a_2 = \sigma^{\mu\nu} \sigma_{\mu\nu}, a_3 = u_\mu \sigma^{\mu\nu} \sigma_{\nu\lambda} u^\lambda. \quad (4)$$

A further possible combination $a_4 = (\epsilon^{\mu\nu\lambda} u_\mu \sigma_{\nu\lambda})^2$ depends on (4),

$$a_4 = 4a_3 + 2a_1a_2. \quad (5)$$

The canonical momenta are

$$P^\mu = -\frac{\partial L}{\partial u_\mu} = -2u^\mu L_1 - 2\sigma^{\mu\alpha} \sigma_{\alpha\beta} u^\beta L_3, \quad (6a)$$

$$S^{\mu\nu} = -\frac{\partial L}{\partial \sigma_{\mu\nu}} = -4\sigma^{\mu\nu} L_2 - 2(\sigma^{\nu\alpha} u^\mu u_\alpha - \sigma^{\mu\alpha} u^\nu u_\alpha) L_3, \quad (6b)$$

where $L_i = \frac{\partial L}{\partial a_i}$. The equations of motion are

$$\dot{P}^\mu = 0, \dot{S}^{\mu\nu} + \sigma^{\mu\lambda} S_\lambda{}^\nu - S^{\mu\lambda} \sigma_\lambda{}^\nu = 0 = \dot{S}^{\mu\nu} + u^\mu P^\nu - P^\mu u^\nu. \quad (7)$$

The conserved Poincare generators are P^μ and $M^{\mu\nu} = S^{\mu\nu} + x^\mu P^\nu - P^\mu x^\nu$. Note that in general the scalars

$$P^\mu P_\mu = M^2 = 4(a_1 L_1^2 + 2a_3 L_1 L_3 - \frac{1}{2} a_2 a_3 L_3^2), \quad (8)$$

$$S^{\mu\nu} S_{\mu\nu} = J^2 = 8(2a_2 L_2^2 + 4a_3 L_2 L_3 - a_1 a_3 L_3^2), \quad (9)$$

$$W^2 = (\epsilon^{\mu\nu\lambda} S_{\mu\nu} P_\lambda)^2 = \frac{1}{2} M^2 J^2 - S^{\mu\nu} P_\nu S_{\mu\alpha} P^\alpha, \quad (10)$$

are operators having non-trivial Poisson Brackets (PB). (10) is the square of the Pauli-Lubanski scalar.

The next task is to specify the constraints such that only a single component of the angular momentum $S^{\mu\nu}$ remains. An appropriate set of constraints is [6]

$$V^\mu \equiv S^{\mu\nu} P_\nu \approx 0. \quad (11)$$

(With this choice W^2 reduces to the standard expression.) Note that only two components of V^μ are independent since $V^\mu P_\mu = 0$, and so they reduce $S^{\mu\nu}$ correctly to a single component, in the anyon frame.. Substituting P^μ and $S^{\mu\nu}$ from (6) in (11), we get

$$V^\mu = 2\sigma^{\mu\nu} u_\nu (4L_1 L_2 - 2a_1 L_1 L_3 - 2a_2 L_2 L_3 - 2a_3 L_3^2) = 0. \quad (12)$$

For the validity of (11), L has to satisfy the differential equation

$$2L_1L_2 - a_1L_1L_3 - a_2L_2L_3 - a_3L_3^2 = 0.$$

The choice of the above condition is not unique and indeed other (interrelated) alternatives are available. Actually these different choices signify distinct frames of reference. The particular choice (11) specifies the so called centre of momentum frame [7,5], where the spatial momentum of the system vanishes. The centre of mass coordinates are a generalization of the non-relativistic definition, with the mass replaced by the energy. This can be further generalized to arbitrary inertial frames (see Pryce in [7]).

Let us now introduce the Lagrangian L , which being a homogeneous function of degree one in the velocities, satisfies

$$u_\mu \frac{\partial L}{\partial u_\mu} + \frac{1}{2} \sigma_{\mu\nu} \frac{\partial L}{\partial \sigma_{\mu\nu}} = L,$$

and is of the generic form

$$L = 2a_1L_1 + 2a_2L_2 + 4a_3L_3, \tag{13}$$

obeying the condition (12).

There are two approaches of deriving the desired Lagrangian. The more fundamental and straightforward one is to solve a consistency condition involving L , M^2 , J^2 and a_i , obtained by eliminating the L_i from (8), (9), (12) and (13). The relation comes in fourth power of L^2 . Each root is a possible choice of L , producing its own RT. It has been shown in [5] that for a relativistic spinning particle, in general M^2 and J^2 can not be fixed arbitrarily. Rather there exists an RT of the form

$$P^2 - f(S^2) = M^2 - f(J^2) = 0. \tag{14}$$

In general the RT should be obtained from (8), which relates M^2 to J^2 , as in (14). (However there is more subtlety involved in the idea of having a non-trivial RT for each L . But we will come back to this point later.) What this means is that even classically, the mass and spin

of a particle can not be fixed indepentently of each other, due to the restriction imposed by the RT. For example, for a particular spin value, (remember that classically the spin can be arbitrary), only specific values of mass of the particle are allowed, which are computed from the RTs and each will have its own L , a root of the consistency equation. Clearly all these models are *inequivalent*. However, these models are somewhat unfamiliar, since in general, their mass and spin are operator valued (see below eqs. (8)-(10)). Unfortunately the consistency equation is very complicated to solve for L .

The alternative and more practicable approach is to proceed via the RT [5]. This means that here one infers an explicit form of an RT, (from other considerations, as for example a linear one as in (16) below), and then tries to construct an L , (by employing the iconal procedure), that agrees with this RT and satisfies the consistency relation as well, (such as the one in (15) below). Generically the L is reconstructed from the RT on the boundary submanifold $a_3 = 0$ and then (12) is used in a generalized eiconal procedure [5] to continue L for non-zero a_3 . An exampe of a non-trivial L is

$$L^2 = \frac{1}{2}[Aa_1 - Ba_2 + ((Aa_1 - Ba_2)^2 - 8ABa_3)^{\frac{1}{2}}], \quad (15)$$

corresponding to the given RT

$$BM^2 - \frac{1}{4}AJ^2 = AB. \quad (16)$$

(The derivation of this L will be given in an enlarged version.)

Let us now compute the canonical PBs between x^μ , P^μ , $\Lambda^{\mu\nu}$ and $S^{\mu\nu}$. Since $\Lambda\Lambda^T = g$ (1), there are only three independent combinations of Λ which we denote by $\phi_i, i = 1, 2, 3$. These ϕ_i 's are taken as canonical coordinates. Let us also define $\sigma^{\mu\nu} = a_i^{\mu\nu}(\phi)\dot{\phi}_i$. The conjugate momenta for ϕ_i are

$$T_i = -\frac{\partial L}{\partial \dot{\phi}_i} = -\frac{1}{2}\left(\frac{\partial \sigma^{\mu\nu}}{\partial \dot{\phi}_i}\right)\left(\frac{\partial L}{\partial \sigma^{\mu\nu}}\right) = \frac{1}{2}a_i^{\mu\nu}(\phi)S_{\mu\nu}. \quad (17)$$

With these definitions a generic PB becomes

$$\begin{aligned}
\{A, B\} &= \left(\frac{\partial A}{\partial x_\mu}\right)\left(\frac{\partial B}{\partial P^\mu}\right) + \left(\frac{\partial A}{\partial \phi_i}\right)\left(\frac{\partial B}{\partial T_i}\right) - \left(\frac{\partial A}{\partial P_i^\mu}\right)\left(\frac{\partial B}{\partial x_\mu}\right) - \left(\frac{\partial A}{\partial T_i}\right)\left(\frac{\partial B}{\partial \phi_i}\right) \\
&= \left(\frac{\partial A}{\partial x_\mu}\right)\left(\frac{\partial B}{\partial P^\mu}\right) - \left(\frac{\partial A}{\partial P^\mu}\right)\left(\frac{\partial B}{\partial x_\mu}\right) + S^{\mu\nu}\left(\frac{\partial A}{\partial S^{\mu\lambda}}\right)\left(\frac{\partial B}{\partial S^{\nu\lambda}}\right) \\
&\quad + \frac{1}{2}\left(\Lambda^\mu{}_\alpha \frac{\partial A}{\partial \Lambda^{\mu\beta}} - \Lambda^\mu{}_\beta \frac{\partial A}{\partial \Lambda^{\mu\alpha}}\right)\frac{\partial B}{\partial S_{\alpha\beta}} - \frac{1}{2}\left(\Lambda^\mu{}_\alpha \frac{\partial B}{\partial \Lambda^{\mu\beta}} - \Lambda^\mu{}_\beta \frac{\partial B}{\partial \Lambda^{\mu\alpha}}\right)\frac{\partial A}{\partial S_{\alpha\beta}}.
\end{aligned} \tag{18}$$

With the help of (18), the PB's are

$$\begin{aligned}
\{P^\mu, x^\nu\} &= -g^{\mu\nu}, \{P^\mu, P^\nu\} = \{x^\mu, x^\nu\} = \{\Lambda^{\mu\nu}, \Lambda^{\alpha\beta}\} = 0 \\
\{\Lambda^{\mu\nu}, S^{\alpha\beta}\} &= \Lambda^{\mu\alpha} g^{\nu\beta} - \Lambda^{\mu\beta} g^{\nu\alpha}, \\
\{S^{\mu\nu}, S^{\alpha\beta}\} &= S^{\mu\alpha} g^{\nu\beta} - S^{\mu\beta} g^{\nu\alpha} + S^{\nu\beta} g^{\mu\alpha} - S^{\nu\alpha} g^{\mu\beta}.
\end{aligned} \tag{19}$$

The Hamiltonian description is pursued by first noting that besides the RT, which is a First Class Constraint (FCC), (any two) of the constraints V^μ , in (12), constitute a Second Class (SC) pair with

$$\{V^\mu, V^\nu\} = S^{\mu\nu} M^2. \tag{20}$$

Consequently $\Lambda^{\mu\nu}$ should also be constrained so that only one angle variable remains. A particularly useful covariant choice is [5]

$$\chi^\mu \equiv \Lambda^{0\mu} - \frac{P^\mu}{M} \approx 0. \tag{21}$$

Again only two of the χ^μ are independent since

$$\chi_\mu(\Lambda^{0\mu} + \frac{P^\mu}{M}) = 0.$$

The total system of four constraints V^1, V^2, χ^1 and χ^2 constitute a SC system and the RT (16) is the only FCC. Thus the Hamiltonian is

$$H = v(\tau)(P^2 - f(S^2)), \tag{22}$$

where $v(\tau)$ is an undetermined multiplier. It is straightforward to compute the intermediate Dirac Brackets (DB). The DB's involving P^μ remain unchanged, whereas x^μ DB's are drastically altered to

$$\{x^\mu, x^\nu\}_I = -\frac{S^{\mu\nu}}{M^2}; \{x^\mu, S^{\nu\lambda}\}_I = \frac{S^{\mu\nu}P^\lambda - S^{\mu\lambda}P^\nu}{M^2}. \quad (23)$$

Actually in the reduced system $S^{01} = \frac{S^{12}P_2}{P^0}$ and $S^{02} = -\frac{S^{12}P_1}{P^0}$, but the $\Lambda^{\mu\nu}$ components are not so simply related to, say Λ^{12} . That is why we prefer to give the algebra in the covariant notation. The final form is reproduced in (26). The Poincare generators P^μ and $M^{\mu\nu}$ DB's are

$$\begin{aligned} \{M^{\mu\nu}, M^{\alpha\beta}\}_I &= M^{\mu\alpha}g^{\nu\beta} - M^{\mu\beta}g^{\nu\alpha} + M^{\nu\beta}g^{\mu\alpha} - M^{\nu\alpha}g^{\mu\beta}, \\ \{M^{\mu\nu}, P^\alpha\}_I &= g^{\mu\alpha}P^\nu - g^{\nu\alpha}P^\mu. \end{aligned} \quad (24)$$

The single FCC, RT is still remaining and v in (24) is still arbitrary. But this is fixed once a scale for x^0 is fixed, *eg.*

$$\Psi = x^0 - t \approx 0. \quad (25)$$

In conjunction with the RT (16), (25) is a SCC system. So the final DB's are

$$\begin{aligned} \{x^\mu, x^\nu\}_{II} &= -\frac{S^{\mu\nu}}{M^2} + \frac{P^\mu S^{0\nu} - P^\nu S^{0\mu}}{M^2 P^0}; \\ \{P^\mu, P^\nu\}_{II} &= \{P^\mu, S^{\alpha\beta}\}_{II} = 0; \{P^\mu, x^\nu\}_{II} = -g^{\mu\nu} + \frac{g^{\mu 0}P^\nu}{P^0}; \\ \{\Lambda^{\mu\nu}, P^\alpha\}_{II} &= \left(\frac{f'}{P^0}\right)g^{0\alpha}\Lambda^\mu{}_\lambda S^{\lambda\nu}; \\ \{x^\mu, S^{\nu\lambda}\}_{II} &= \frac{S^{\mu\nu}P^\lambda - S^{\mu\lambda}P^\nu}{M^2} - \left(\frac{S^{0\nu}P^\lambda - S^{0\lambda}P^\nu}{M^2 P^0}\right)P^\mu; \\ \{S^{\mu\nu}, S^{\alpha\beta}\}_{II} &= S^{\mu\alpha}(g^{\nu\beta} - \frac{P^\nu P^\beta}{M^2}) - S^{\mu\beta}(g^{\nu\alpha} - \frac{P^\nu P^\alpha}{M^2}) \\ &\quad + S^{\nu\beta}(g^{\mu\alpha} - \frac{P^\mu P^\alpha}{M^2}) - S^{\nu\alpha}(g^{\mu\beta} - \frac{P^\mu P^\beta}{M^2}), \end{aligned} \quad (26)$$

where $f' = \frac{\partial f}{\partial J^2}$, with f obtained from (14). But here the $S^{\mu\nu}$ algebra is simple since

$$\{S^{12}, S^{12}\} = 0; \{x^i, S^{12}\} = -\frac{S^{12}P^i}{M^2}.$$

All the $\Lambda^{\mu\nu}$ DB's are not shown, but it can be seen, (see *eg.* $\{\Lambda^{\mu\nu}, P^\alpha\}_{II}$ in (26)), that only the *right* index of $\Lambda^{\mu\nu}$ transform under Poincare generators. The Hamiltonian

$$H = P^0 = [P^2 + f(J^2)]^{\frac{1}{2}}, \quad (27)$$

generates the time translations

$$\frac{dA}{dt} = \frac{\partial A}{\partial t} + \{H, A\}_{II}.$$

The final Poincre DB algebra is the same as in (24).

However, due to the presence of nonzero DB between x^μ 's and the mixed DB's between position and angle variables, it is better to switch over to the Pryce-Newton-Wigner variables [5,7], in terms of which the DB's simplify and the variables are amenable to quantization, such that the DB's are replaced by commutators.

Now we shall discuss the special type of Lagrangian which was mentioned before, which does not obey the RT restriction. The following Lagrangian

$$\mathcal{L}^2 = [a_1 M^2 + \frac{1}{2} a_2 J^2 + 2MJ(\frac{a_1 a_2}{2} + a_3)^{\frac{1}{2}}], \quad (28)$$

with M and J as *arbitrary numerical* parameters, also satisfies the constraint relation (12) with

$$P^\mu P_\mu = M^2, S^{\mu\nu} S_{\mu\nu} = J^2, W = MJ. \quad (29)$$

Thus in this particular case, P^2 and S^2 can be numbers which are independently fixed. (Note that in the 3+1-dimensional case in [5], this type of solution appeared only in the limit of vanishing a_4 , whereas here since a_4 of (5) is not present from the very beginning, this is also a solution without any approximation). If one were able to solve explicitly

the consistency condition mentioned before, (15) and (28) should appear as two of the independent roots. This is the subtlety we mentioned before and for reasons obvious by now, we believe (28) should be chosen to describe an anyon.

The constraint structure of this $\mathcal{L}$ in (28) is essentially different since it has two independent FC's [2],

$$\Phi_1 = P^\mu P_\mu - M^2, \Phi_2 = \epsilon^{\mu\nu\lambda} S_{\mu\nu} P_\lambda - MJ. \quad (30)$$

(Due to the relation (10), S^2 is not independent.) The Φ 's are the two Casimir operators of the 2+1-dimensional Poincare algebra. The Hamiltonian is

$$H = v_1 \Phi_1 + v_2 \Phi_2.$$

Therefore apart from fixing the time scale to gauge fix the mass shell condition, (as in (15)), we have to fix another gauge for Φ_2 , which we can take as χ^1 of (23). We still require two more constraints which are as in (12). Interestingly the DB's due to the SCC's Φ_2 , χ^1 , V^1 and V^2 remain the same as in (25). The reduced H is $v_1 \Phi_1$. The final H is

$$H = P^0 = [P^2 + M^2]^{\frac{1}{2}}$$

.

Finally we come back to the model proposed in [3]. In order to forge a comparison between our results and those of [3], the following identifications [8] are made,

$$\sigma^{\alpha\beta} \equiv \frac{1}{2}(n^\alpha \dot{n}^\beta - \dot{n}^\alpha n^\beta), S^{\alpha\beta} \equiv (n^\alpha p_n^\beta - p_n^\alpha n^\beta), \quad (31)$$

where $n^2 = -1$ and $p_n^\mu = -\frac{\partial L}{\partial \dot{n}^\mu}$. Even though the DB structure of [3] and ours agree, the Lagrangians are different. Our expression in the new variables become

$$\bar{\mathcal{L}}^2 = M^2 \dot{x}^2 - \frac{J^2 \dot{n}^2}{4} + MJ \epsilon^{\mu\nu\alpha} n_\mu \dot{n}_\nu \dot{x}_\alpha, \quad (32)$$

whereas in [3], the expression is

$$\tilde{L}^2 = m^2 \dot{x}^2 - \alpha^2 \dot{n}^2 + 2m\alpha \epsilon^{\mu\nu\alpha} n_\mu \dot{n}_\nu \dot{x}_\alpha + m^2 (\dot{x}^\mu n_\mu)^2. \quad (33)$$

Notice that a term of the form of the last one in $\tilde{L}$ in (33) is missing from $\bar{\mathcal{L}}$ in (32). This casts some doubt on the validity of the Lagrangian in [3] because, (although naively the Poincare algebra is satisfied), complications are involved in the transformation properties. For example, as we mentioned before, only the right index of $\Lambda^{\mu\nu}$ transform under Poincare generators, whereas no such distinction is discernable in [3]. Furthermore, the constraint analysis of [3] is very obscure. The authors have started with a simple Lagrangian for the free anyon, where the spin parameter does not appear in the Lagrangian. They then assume that the constraints of this system will be preserved in the more detailed Lagrangian, where the spin value explicitly appears. The adhoc nature of introducing the Pauli-Lubanski constraint is questionable. All these factors are probably responsible for the mismatch between the Lagrangian in [3], (obtained in a heuristic fashion), and our expression.

To conclude, we have derived two possible expressions for the Lagrangian of a free anyon. One of them obeys the Regge Trajectory restriction between its mass and spin values, whereas the other does not. But in the latter model, the anyon mass and spin are *independent c-number parameters*. Obviously this candidate Lagrangian is a better choice to describe an anyon, instead of the one obeying RT and having operator valued mass and spin. A detailed Hamiltonian analysis for both of these models is carried through and all the relevant Dirac Brackets are computed. The variables suitable for quantization are also discussed. Finally differences between our model and a recently proposed model [3] are pointed out. The problems of introducing interactions and a non-trivial gravitational background are under study.

Acknowledgements: I thank Dr. P. Mitra and Narayan Banerjee for discussions.

References:

- [1] F.Wilczek, *Fractional Statistics and Anyon Superconductivity*, (World Scientific, 1990)
- [2] R.Jackiw and V.P.Nair, Phys.Rev.**D43** (1991)1933
- [3] M.Chaichian, R.Gonzalez Felipe and D.Loiz Martinez, Phys.Rev.Lett. **71** (1993)3405
- [4] P.A.M.Dirac, *Lectures on Quantum Mechanics*, (Yeshiva University Press, New York, 1964)
- [5] A.J.Hanson and T. Regge, Ann.Phys.(N.Y.) **87** (1974)498; see also A.J.Hanson, T. Regge and C. Teitelboim, *Constrained Hamiltonian Systems*, (Roma, Academia Nazionale dei Lincei, 1976)
- [6] J.Wessynhoff and A.Raabe, Acta Physica Polonica **9** (1947)7; J.Wessynhoff, *ibid* (1947)26,34; J.L.Synge, *Relativity: the Special Theory*, (North Holland, Amsterdam, 1955)
- [7] M.H.L.Pryce, Proc.Roy.Soc. London, **Ser A150** (1935)166; *ibid* **195** (1948)62
- [8] A. O. Barut, *Electrodynamics and the Classical Theory of Fields and Particles*, (Macmillan Co., New York, 1964)